

Problem 25 (From the lecture)

Verify the following statements from the lecture:

- (1) The actions of S_n and a group G on $V^{\otimes n}$ as defined in the lecture commute.
- (2) The tensor coefficients in the irrep corresponding to the Young tableau

1	3
2	

satisfy $a_{ijk} + a_{jki} + a_{kij} = 0$ and $a_{ijk} = -a_{jik}$.

(Optional) Problem 26 (Irreps of SU(N))

As discussed in the lecture, the Schur-Weyl duality dictates that distinct irreps of SU(N) are given by different Young diagrams with at most N rows. Using the Young diagrams, the decomposition of tensor product of irreps $T_1 \otimes T_2$ can be computed in a pictorial way as described below:

- (1) assign distinct labels to boxes in each row of T_2 , e.g.

1	1
2	
- (2) attach boxes of T_2 row by row to T_1 in all possible ways that result in another semistandard Young tableau.
- (3) after attaching each row, one should keep only those tableaux that when reading *from right to left while going from top to bottom*, the sequence of numbers attached should contain at least as many 1's as 2's, 2's as 3's, etc at any point.

Below is an example:

$$\begin{array}{c}
 \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \otimes \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \cong \begin{array}{|c|c|c|c|} \hline & & 1 & 1 \\ \hline & 2 & & \\ \hline \end{array} \oplus \begin{array}{|c|c|c|c|} \hline & & 1 & 1 \\ \hline & & & 2 \\ \hline \end{array} \oplus \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 1 & 2 \\ \hline \end{array} \\
 \\
 \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 1 & \\ \hline 2 & & \end{array} \oplus \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 2 & \\ \hline 1 & & \end{array} \oplus \begin{array}{|c|c|c|c|} \hline & & & 1 \\ \hline & & 1 & \\ \hline & 1 & 2 & \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline & \\ \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline & \\ \hline & 1 \\ \hline \end{array} \quad (1)
 \end{array}$$

Note that the tableau

		1	2
	1		

 is not allowed because it violates condition (3).

- (1) Perform the decomposition for the following tensor product for SU(3):

$$\begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \otimes \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array}$$

- (2) The dimension of the irrep for $SU(N)$ is given by $D = \prod_b \frac{(N+d(b))}{h(b)}$, where the multiplication goes over all boxes b in the diagram. $h(b)$ is the hook length of box b , and $d(b)$ defined in the following example diagram

0	1	2	3
-1	0	1	2
-2	-1	0	1

Check that the dimensions match in the decomposition you just performed. For more details consult textbooks, e.g. Jones, *Groups, representations and physics*, 2nd Ed., IoP 1998.