

P 24. $D_4 = \langle r, s \mid r^4 = s^2 = (rs)^2 = 1 \rangle$

$$= \{ 1, r, r^2, r^3, s, rs, r^2s, r^3s \}$$

$$\left(\begin{matrix} \parallel & \parallel & \parallel \\ sr^3 & sr^2 & sr \end{matrix} \right)$$

(1) $D_4 = \{ 1 \} \cup \{ r, r^3 \} \cup \{ r^2 \} \cup \{ s, r^2s \} \cup \{ rs, r^3s \}$

$$=: [1] \cup [r] \cup [r^2] \cup [s] \cup [rs]$$

$$C_1 = 1; \quad C_2 = r + r^3; \quad C_3 = r^2; \quad C_4 = s + r^2s; \quad C_5 = rs + r^3s$$

(2)

| | C_1 | C_2 | C_3 | C_4 | C_5 |
|-------|-------|---------------|-------|---------------|---------------|
| C_1 | C_1 | C_2 | C_3 | C_4 | C_5 |
| C_2 | | $2C_1 + 2C_3$ | C_2 | $2C_5$ | $2C_4$ |
| C_3 | | | C_1 | C_4 | C_5 |
| C_4 | | | | $2C_1 + 2C_3$ | $2C_2$ |
| C_5 | | | | | $2C_1 + 2C_3$ |

(3) $L_{jk} = \sum_i C_{ij}^k y^i$

$$L = \begin{pmatrix} y^1 & y^2 & y^3 & y^4 & y^5 \\ 2y^2 & y^1 + y^3 & 2y^2 & 2y^5 & 2y^4 \\ y^3 & y^2 & y^1 & y^4 & y^5 \\ 2y^4 & 2y^5 & 2y^4 & y^1 + y^3 & 2y^2 \\ 2y^5 & 2y^4 & 2y^5 & 2y^1 & y^1 + y^3 \end{pmatrix}$$

$$\lambda_a = y^1 - y^3 \quad m_1 = 1$$

$$\lambda_b = y^1 + 2y^2 + y^3 - 2y^4 - 2y^5 \quad m_2 = 2$$

$$\lambda_c = y^1 - 2y^2 + y^3 + 2y^4 - 2y^5 \quad m_3 = 1$$

$$\lambda_d = y^1 - 2y^2 + y^3 - 2y^4 + 2y^5 \quad m_4 = 2$$

$$\lambda_e = y^1 + 2y^2 + y^3 + 2y^4 + 2y^5 \quad m_5 = 2$$

②

$$\lambda_\mu = \frac{1}{n_\mu} \sum_{i=1}^r m_i \chi_\mu([C_i]) y^i \Rightarrow$$

$$\chi_a = n_a (1, 0, -1, 0, 0)$$

$$\chi_b = n_b (1, 1, 1, -1, -1)$$

$$\chi_c = n_c (1, -1, 1, 1, -1)$$

$$\chi_d = n_d (1, -1, 1, -1, 1)$$

$$\chi_e = n_e (1, 1, 1, 1, 1)$$

$$n_\mu = \left[\frac{(g)}{\sum_{i=1}^r m_i \left| \frac{\chi_\mu([C_i])}{n_\mu} \right|^2} \right]^{\frac{1}{2}} \Rightarrow \begin{aligned} n_a &= 2 \\ n_b &= n_c = n_d = n_e = 1 \end{aligned}$$

identify $[r] = C_4(\theta)$, $[r^2] = C_2(\theta)$, $[s] = C_2'$, $[rs] = C_2''$

We recover the character table:

Character table for point group D_4

| D_4 | E | $2C_4(z)$ | $C_2(z)$ | $2C_2'$ | $2C_2''$ | linear functions, rotations | quadratic functions | cubic functions |
|-------|----|-----------|----------|---------|----------|-----------------------------|---------------------|--|
| A_1 | +1 | +1 | +1 | +1 | +1 | - | x^2+y^2, z^2 | - |
| A_2 | +1 | +1 | +1 | -1 | -1 | z, R_z | - | $z^3, z(x^2+y^2)$ |
| B_1 | +1 | -1 | +1 | +1 | -1 | - | x^2-y^2 | xyz |
| B_2 | +1 | -1 | +1 | -1 | +1 | - | xy | $z(x^2-y^2)$ |
| E | +2 | 0 | -2 | 0 | 0 | $(x, y) (R_x, R_y)$ | (xz, yz) | $(xz^2, yz^2) (xy^2, x^2y) (x^3, y^3)$ |

<http://symmetry.jacobs-university.de/cgi-bin/group.cgi?group=304&option=4>

