


P 28. use $\chi_{v \otimes v_2} = \chi_{v_1} \cdot \chi_{v_2}$ and the character theory.

P 29.



$D_3 \cong S_3$ three conj. classes

$$E \leftrightarrow [()]$$

$$C_3 \leftrightarrow [(123)]$$

$$C_2' \leftrightarrow [(12)]$$

$$\chi_v(E) = 6$$

$$\chi_v(C_3) = 0 \quad (\because 1, 2, 3 \text{ all switch places, all } 0 \text{ on diagonal})$$

$$\chi_v(C_2') = 0 \quad (\sigma: \delta y_3 \rightarrow \delta y_3, \delta x_3 \rightarrow -\delta x_3)$$

$$n_p = \langle \chi_p, \chi_v \rangle$$

$$n_{A_1} = \frac{1}{6} 1 \times 6 = 1$$


$$n_{A_2} = \frac{1}{6} 1 \times 6 = 1$$

$$n_E = \frac{1}{6} 2 \times 6 = 2$$


$$V \cong A_1 \oplus A_2 \oplus 2E$$

	E	2C ₃	3C ₂ '
A ₁	1	1	1
A ₂	1	1	-1
E	2	-1	0

$$v: 6 \quad 0 \quad 0$$



C_3
 \rightarrow



$$\delta x, \delta y, \delta x, \delta y, \delta x, \delta y$$

$$(1, 0, 0, 0, 0, 0) \rightarrow$$

$$(0, 0, -\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, 0)$$

$$M(C_3) = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{2} & -\frac{\sqrt{3}}{2} & \dots & 0 \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} & 0 & \dots & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}$$