

P 16.  $D = \{ \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}, z = e^{i\theta} \in U(1) \}$

(a)  $d \in D \quad u = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \in SU(2) \quad |\alpha|^2 + |\beta|^2 = 1$

$$udu^{-1} = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & -\beta \\ \bar{\beta} & \alpha \end{pmatrix}$$

$$= \begin{pmatrix} |\beta|^2 z + |\alpha|^2 \bar{z} & \alpha\beta (-z + \bar{z}) \\ \bar{\alpha}\bar{\beta} (-z + \bar{z}) & |\alpha|^2 \bar{z} + |\beta|^2 z \end{pmatrix} \in D$$

$$\Rightarrow \alpha = 0 \text{ or } \beta = 0$$

$$N_{SU(2)}(D) = \{ \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}, z \in U(1), \cup \begin{pmatrix} 0 & -\bar{z} \\ z & 0 \end{pmatrix}, z \in U(1) \}$$

$$= D \cup \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} D$$

(b)  $N_{SU(2)}(D)/D = \{ \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} D = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} D,$

$$\begin{pmatrix} 0 & -\bar{z} \\ z & 0 \end{pmatrix} D = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} D \quad \{ \cong \mathbb{Z}_2$$

(c)  $\begin{pmatrix} \alpha & 0 \\ 0 & \bar{\alpha} \end{pmatrix} \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \begin{pmatrix} \bar{\alpha} & 0 \\ 0 & \alpha \end{pmatrix} = \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix}$

$$\begin{pmatrix} 0 & -\bar{z} \\ z & 0 \end{pmatrix} \begin{pmatrix} z & 0 \\ 0 & \bar{z} \end{pmatrix} \begin{pmatrix} 0 & \bar{z} \\ -z & 0 \end{pmatrix} = \begin{pmatrix} \bar{z} & 0 \\ 0 & z \end{pmatrix}$$

(d) should at least contain  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . and some

$$a = \begin{pmatrix} 0 & z \\ -\bar{z} & 0 \end{pmatrix} \text{. then it contains } a^2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\& \quad a^3 = \begin{pmatrix} 0 & -z \\ \bar{z} & 0 \end{pmatrix} . \quad \text{it's not isomorphic to } \mathbb{Z}_2.$$

(2)

NB ( $N_{SU(2)}(D)/D$ ) is not a subgroup of  $SU(2)$   
or  $N_{SU(2)}(D)$

P 17.

$G$ -set  $X$ .  $\phi: G \rightarrow S_X$

(a) effective  $\Leftrightarrow \phi$  injective, i.e.  $\phi(g) = 1$  iff  $g=1$

$\forall g \neq 1$ .  $\exists x$  s.t.  $gx_1 = x_2 \neq x_1 \Leftrightarrow \forall g \neq 1$ .  $\phi(g)$  is a nontrivial  
permutation  
 $\phi(g) \neq 1$

(b)  $\{f_i\}$  are ineffective  $\forall f \in G$

$$f_i \circ f_j x = f_i \cdot f_j x' = x' \quad (\forall x \in X)$$

$$f_j \circ f_i x = f_j x = x'$$

$$\Rightarrow f_i \circ f_j = f_j \circ f_i \quad \forall f \in G$$

trivial to show  $\{f_i\}$  is a group

$$\Rightarrow H = \{f_i : f_i x = x \ \forall x \in X\} \triangleleft G$$

(c) define the action  $G/H \times X \rightarrow X$

$$(fH)x := fx$$

$$\forall x \in X, \text{ s.t. } (fH)x = x \Leftrightarrow fx = x \Leftrightarrow f \in H \Leftrightarrow fH = H = 1_{G/H}$$

P(8).  $X$  a finite  $G$  set.

$G$ -action transitive  $\Rightarrow$  one orbit  $= X$

Burnside's lemma  $\Rightarrow |G| = \sum_{g \in G} |X^g|$

If all  $g$ 's have fixed points.  $\sum_{g \in G} |X^g| \geq \sum_{g \in G} 1 = |G|$   
 Equality holds iff  $\forall g. |X^g| = 1$

But  $|X^e| = |X| > 1$

$\Rightarrow |X^g| = 1$  for some  $g$ .